



WSD COMMAND POUR ON LOUSICIDE FOR SHEEP

Along with its Australian made drench range, WSD produces various products for controlling Lice. **WSD Command** is the flagship of the Lice Control range. It is a part of the Insect Growth Regulator (IGR) group of products.

WSD Command attacks immature Lice before they hatch, controlling the populations for up to 20 weeks.

WSD Command is fully registered with the Australian Pesticides and Veterinary Medicines Authority (APVMA) - the peak body for animal health products in Australia.

PLEASE NOTE:

Information correct at time of printing. No warranty expressed or implied is given with the sale of WSD products. Use of these products is the responsibility of the owner of the animal(s) being treated. It is recommended that you contact your local veterinarian before using any product. Product efficacy will vary between properties. WSD suggests that good management practices are followed to ensure resistance to any chemical is kept minimal. WSD cannot guarantee any product will work to one hundred percent effectiveness.

Correct application

When using any product it is important you read and understand the label directions for use.

WSD Command is an IGR chemical. The effectiveness of treatment by any IGR depends on the chemical coming into direct contact with lice.

Sheep **MUST** be clean shorn with no residual clumps or tufts of wool. Animals with extensive skin folds or wrinkles are unlikely to be treated effectively by **WSD Command**. It may be necessary to cull sheep with extensive wrinkles before using **WSD Command**.

Animals affected by lumpy wool (mycotic dermatitis) must be treated for the condition and clean shorn before applying **WSD Command**.

There may be live lice on sheep for a considerable period following treatment. Lice populations may not be controlled for up to 12 weeks after application. Infested sheep should be separated from clean animals during the treatment period. The effectiveness of **WSD Command** may be reduced if applied during extreme climatic conditions, or if animals are under severe nutritional stress.

WSD Command must be applied within 7 days after shearing. For optimum results, apply within 12 hours of shearing.

Application equipment must be suitable for this type of product. Contact WSD if you need specialist equipment. Equipment must be calibrated immediately prior to and regularly during use, to ensure that the correct dose is administered.

Protective clothing specified on the label, must be worn, including by anyone handling the sheep soon after treatment.

WSD Command

WSD Command is an Insect Growth Regulator

Applied as Pour On application

Contains scourable dye, 66 days ESI & 14 days meat withholding

Available in 20L drum & 5L backpack

Dose Rate 5 ml per 10Kg body weight

