



## SPECIFICATION SHEET

# WSD CY-GUARD SPRAY ON

### Product Description

WSD CY-GUARD SPRAY ON is an external application for preventing blowfly strike in sheep and lambs.

### Type of Preparation

WSD CY-GUARD SPRAY ON is a ready-to-use formulation containing 60g/L **Cyromazine** and a scourable dye for topical application.

### Indications

**Sheep and lambs:** Prevention for up to 11 weeks but NOT treatment of blowfly strike by *Lucilia cuprina* in sheep.

**Milking sheep:** Should not be used in sheep producing milk for human consumption.

### Dosage

Always refer to the label.

Always apply WSD CY-GUARD SPRAY ON undiluted from the original container.

It is important to use an approved applicator to deliver a band onto the fleece close to the areas to be protected. There are different application rates and patterns for protection against poll, body and/or crutch strike, and rates also depend on wool length.

If a second treatment is required during a season, it should be applied less than 11 weeks after the first treatment with WSD CY-GUARD SPRAY ON or less than 14 weeks after treatment with WSD CY-GUARD SC.

### Advice on use

- WSD CY-GUARD SPRAY ON is convenient to use but may not provide protection for as long as jetting or spraying with WSD CY-GUARD SC.
- WSD CY-GUARD SPRAY ON can be used to achieve moderate periods of fly-strike prevention of up to 11 weeks, including where it is not possible to correctly apply WSD CY-GUARD SC by dipping or jetting.
- Duration of activity and effectiveness of WSD CY-GUARD SPRAY ON depend on correct application technique.
- WSD CY-GUARD SPRAY ON must not be diluted before use.
- It is IMPORTANT to use WSD CY-GUARD SPRAY ON BEFORE the anticipated onset of a major fly wave. This may be at a particular time of year OR following unseasonal rain or if exceptional blow-fly activity is detected.
- WSD CY-GUARD SPRAY ON may be used in sheep with less than 6 weeks wool.
- Consider using WSD CY-GUARD SPRAY ON for sheep severely affected by grass seeds. WSD CY-GUARD SPRAY ON is also suitable for use in sheep with open wounds provided it is not applied directly onto the wounds.
- Do NOT treat wet sheep. Heavy rain soon after the application of this product may also reduce the period of protection, so avoid use if heavy showers are forecast.
- This product must NOT be mixed with any other product.

### Withholding periods

**Meat withholding:** 7 days

**Export slaughter interval:** 14 days

**Milk withholding:** Not for use in sheep producing milk for human consumption

**Wool withholding:** 60 days (2 months).

### Shelf Life

24 months from the date of manufacture

### Sizes Available

20 litre Drum



**DIRECTIONS FOR USE:**  
 Restraints: DO NOT USE on sheep with less than 6 weeks wool.  
 DO NOT USE on sheep which are producing or may in the future produce milk or milk products for human consumption.  
 Contraindications: This product should not be used on open wounds, e.g. those resulting from marking or milking operations.  
 This product should not be applied during gusty or windy weather as spray drift will reduce the period of protection.

# WSD CY-GUARD SPRAY ON

### WITHHOLDING PERIODS:


**MEAT:** DO NOT USE less than 7 days before slaughter for human consumption.  
**MILK:** DO NOT USE on sheep which are producing or may in the future produce milk or milk products for human consumption.  
**WOOL:** DO NOT USE less than 2 months before shearing or fibre collection.

### TRADE ADVICE:

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 14 days before slaughter for export.

### CRITICAL COMMENTS

MONTHS SINCE SHEARING	TOTAL VOLUME (ml)	VOLUME PER TREATMENT BAND (ml)	NO. OF TREATED BANDS	SHEEP PER 1L	CRITICAL COMMENTS
Up to 6	34	17	2	29	1. Use a WSD recommended Spray-On applicator gun. 2. Ensure the applicator gun is clean.
7	38	19	2	26	3. Cartridge applicator gun using an accurate volume measure.
8	44	22	2	22	4. Ensure all areas to be protected are treated.
9	50	25	2	20	5. When preventing body strike ensure bands overlap along the midline.
10	56	28	2	17	6. This product is not ideal for treating existing strikes. Treated larvae may take 3-4 days to die. Use a product such as WSD Diquat as a dressing for existing strikes.
Lambs less than 15kg	17	17	1	58	7. Heavy rain following the application of this product could diminish the period of protection.
FOR THE PREVENTION OF POLL STRIKE:					
Up to 6	51	17	3	19	8. WSD CY-GUARD SPRAY is coloured for band identification. The scourable dye will disappear shortly after treatment.
7	57	19	3	17	SEE DIAGRAM
8	66	22	3	15	
9	75	25	3	13	
10	84	28	3	11	
Lambs less than 15 kg	34	17	2	29	



FOR THE PREVENTION OF POLL STRIKE: Polls and sheep susceptible to poll strike should be treated. Apply at least 5 mL of WSD CY-GUARD SPRAY to the areas requiring protection, around the poll and horns.  
 NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

### GENERAL INSTRUCTIONS:

DO NOT DILUTE. DO NOT mix with any other product. Apply directly from the container using WSD recommended Spray-on applicator gun. Except for small lambs under 15kg, the dose volume is determined by the number of months since shearing. Apply WSD CY-GUARD SPRAY according to the diagrams and tables in the Directions for Use. WSD CY-GUARD SPRAY may be applied to all breeds of sheep, providing protection against flystrike. WSD CY-GUARD SPRAY is applied as a low-volume treatment onto or near areas where protection is required. When applying, the nozzle of the applicator gun should be approximately 25cm above the head, to achieve a 15cm wide band at each pass.

### SAFETY DIRECTIONS:

Will irritate the eyes and skin. Avoid contact with eyes and skin. When using the product wear rubber gloves. Wash hands after use.

### FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

### PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT:

DO NOT contaminate dams, rivers or streams with the product or used containers.

### MATERIAL SAFETY DATA SHEET:

If additional hazard information is required, refer to the Material safety data sheet. For a copy, phone WSD (08) 9321 2888 or visit [www.westernstock.com.au](http://www.westernstock.com.au)

### DISPOSAL:

Triple or preferably pressure rinse containers and dispose of rinsate in a disposal pit as described below. DO NOT dispose of undiluted chemicals on-site. Break, crush or puncture and bury empty containers in a local authority landfill. If not available bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roads. Empty containers and product should not be burned.

### STORAGE:

Store below 30°C (room temperature). Store in original container tightly closed in a dry, cool place. DO NOT store in direct sunlight. DO NOT re-use the container. Keep out of reach of children.

### WARRANTY AND EXCLUSION OF LIABILITY:

This product is warranted for the purpose specifically recommended by Western Stock Distributors Pty Ltd when used strictly as directed on this label. All other warranties and obligations for livestock, whether expressed or implied by statute or otherwise, are excluded to the full extent that exclusion is permitted by law.

APVMA Approval No. 61663/20/L1107

Batch No:

Expiry: