



## WSD DIAZINON

### Product Description

WSD DIAZINON is a treatment for external parasites of sheep, cattle, goats and pigs.

### Type of Preparation

WSD DIAZINON is an emulsifiable concentrate containing 200g/L Diazinon- an organophosphate insecticide- to be diluted with water and used as a spray or dressing.

### Indications

**Sheep and lambs:** Blowfly dressing for struck sheep and around marking wounds on lambs.

**SPECIAL NOTE:** The use of diazinon to treat sheep lice is not permitted except under a permit issued by the APVMA.

**Milking sheep:** Should not be used in sheep producing milk for human consumption.

**Cattle:** Biting and sucking lice (*Bovicola bovis*, *Linognathus vituli*, *Haematopinus eurysternus*).

**Dairy Cattle:** Suitable for cattle producing milk for human consumption (see withholding period information)

**Goats:** Lice (*Damalinea caprae*)

**Dairy Goats:** Suitable for goats producing milk for human consumption (see withholding period information)

**Pigs:** Lice (*Haematopinus suis*) and mange (*Sarcoptes scabiei*)

### Dosage

Always refer to the label.

**Sheep:** Blowfly treatment- Dilute 5ml/L (1g/L) in water and saturate the affected area. Use a brush or suitable applicator to avoid contact by the operator.

Marking wounds- Dilute 5ml/2L (0.5g/L) in water and apply to the wool around the wound.

**Goats:** Lice- Hand spray: 100ml/40L (0.5g/L) in water. Thoroughly wet goats with spray. A second spraying may be necessary 16 to 17 days later.

**Cattle:** Lice- Hand spray: 1L/400L (0.5g/L) in water. Spray thoroughly using 5 to 10 L per head. A second spraying 16 to 17 days later is necessary to break the life cycle of the lice.

**Pigs:** Lice, mange- Hand spray: 100ml/40L (0.5g/L) in water. Thoroughly wet pigs with spray. Use three applications at 10 day intervals.

### Advice on use

- WSD DIAZINON must never be applied undiluted and should be correctly diluted before use.
- All preparations containing diazinon should be treated with CAUTION. Personal protective equipment is recommended for operators when diluting or using diazinon preparations.
- WSD DIAZINON must not be stored in any container other than the original.
- There is widespread resistance to diazinon among populations of the sheep blowfly *Lucilia cuprina*. Animals treated with diazinon alone may still be susceptible to re-strike.
- Diazinon is no longer permitted to be used for dipping or jetting of sheep for the treatment of lice. WSD provides a range of alternative products for the treatment or control of lice including WSD FLOCKMASTER MKII or WSD COMMAND.

### Withholding periods

#### Meat withholding:

Sheep, goats and pigs: 14 days

Cattle: 3 days

Export slaughter interval: 3 days (cattle) and 21 days (sheep)

#### Milk withholding:

Sheep: Not for use in sheep producing milk for human consumption

Cattle: 6 hours

Goats: 48 hours


Wool withholding: 30 days.

#### Shelf Life

24 months from the date of manufacture

#### Sizes Available

5 litre Drum, 20 litre Drum



## WSD DIAZINON

FOR SHEEP, CATTLE, GOATS AND PIGS

### DIRECTIONS FOR USE

#### RESTRAINTS:

DO NOT use this product for dipping or jetting sheep.

DO NOT USE on female sheep which are producing or may in the future produce milk or milk products for human consumption.

Animal/Pest	Treatment	Rate	Critical Comments
SHEEP Blowfly <i>Lucilia cuprina</i>	Dressing	5mL per 1L of water (1.0g/L Diazinon)	Saturate struck area thoroughly. Use brush on affected area.
CATTLE Lice <i>Linognathus vituli</i> <i>Damalinea bovis</i> <i>Haematopinus eurysternus</i>	Spray	1L per 400L of water (0.5g/L Diazinon)	Use 5-10L per head. Spray thoroughly and repeat spray 15 days later to ensure that lice life cycle is broken.
GOATS Lice <i>Damalinea caprae</i>	Spray	100ml per 40L of water (0.5g/L Diazinon)	Spray goats and repeat spray 15 days later to ensure that lice life cycle is broken.
PIGS Lice <i>Haematopinus suis</i> Mange <i>Sarcoptes sp.</i>	Spray	100ml per 40L of water (0.5g/L Diazinon)	Spray pigs thoroughly. Repeat spray at 10 day intervals three times for best effects.

**NOT TO BE USED FOR ANY OTHER PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

### WITHHOLDING PERIODS:

MEAT (Sheep, Pigs, and Goats): DO NOT USE less than fourteen (14) days before slaughter for human consumption.

MEAT (Cattle): DO NOT USE less than three (3) days before slaughter for human consumption.

MILK (Sheep): DO NOT USE on female sheep which are producing or may in the future produce milk or milk products for human consumption.

MILK (Goats): Milk collected from does within forty-eight (48) hours following treatment MUST NOT BE USED for human consumption or processing. This milk should not be fed to kids.

MILK (Cattle): Milk collected from cows within six (6) hours following treatment MUST NOT BE USED for human consumption or processing. This milk should not be fed to bobby calves.

WOOL: DO NOT USE less than 1 month before shearing or fibre collection.