



WSD FENBENDAZOLE 100

Product Description

WSD FENBENDAZOLE 100 is a broad spectrum treatment for roundworm and tapeworm parasites of cattle and horses

Type of Preparation

WSD FENBENDAZOLE 100 is a suspension for oral use as a drench containing 100g/L Fenbendazole- a benzimidazole (BZ).

Indications

Cattle:

Mature and immature stages of the following worms.

Barber's pole worm (*Haemonchus spp*), hair worm or small hair worm (*Trichostrongylus spp*), small brown stomach worm (*Ostertagia ostertagi* including hypobiotic larvae), small intestinal worm (*Cooperia*), whipworm (*Trichuris*), lungworm (*Dictyocaulus viviparus*), *Capillaria spp*, hookworm (*Bunostomum sp*). Fenbendazole aids in the control of tapeworms (*Moniezia spp*).

Dairy Cattle: Suitable for cattle producing milk for human consumption (see withholding period information)

Horses:

Mature and immature large strongyles (*Strongylus spp*), small strongyles (*Cyathostomidae*), ascarids (*Parascaris equorum*), pinworms (*Oxyuris equi*), *Strongyloides westeri*.

Dosage

Always refer to the label. BZ's generally have low toxicity but to avoid residue problems, it is good practice to separate into different weight classes if there is a large variation in age and weight.

Cattle: Dose at the rate of 3.75ml per 50kg body weight.

Horses: Dose at the rate of 1ml per 10kg body weight

Advice on use

- WSD FENBENDAZOLE 100 is a reliable broad-spectrum oral drench for cattle and horses of all ages.
- WSD FENBENDAZOLE 100 is safe for use in pregnant mares and stud males.
- It is effective in treating severe scouring due to *Ostertagia*-type II ostertagiosis in adult cattle.
- It is often best administered to adult cattle with a drenching hook.
- The repeated use of any drench is likely to result in the emergence of resistance among cattle parasites. WSD provides a range of treatments for cattle parasites and consideration should be given to utilizing more than one treatment group in a worm control program.

Withholding periods

Meat withholding: 14 days

Export slaughter interval: 14 days

Milk withholding: Nil. However, it is generally recommended when treating dairy herds to treat one third of the herd each day over a successive 3 day period.

Shelf Life

24 months from the date of manufacture

Sizes Available

1 litre Twin-Neck, 5 litre Backpack or Quad, 10 litre Drum



WSD FENBENDAZOLE 100

ORAL ANTHELMINTIC FOR CATTLE AND HORSES

Containing Fenbendazole, a member of the benzimidazole family of drenches. It is effective in cattle and horses against benzimidazole sensitive strains of the following internal parasites:

CATTLE

Haemonchus placei
Ostertagia spp
Trichostrongylus spp
Oesophagostomum radiatum
Nematodirus spp
Cooperia spp
Dictyocaulus viviparus
Trichuris spp
Capillaria spp

HORSES

S. vulgaris, *S. edentatus*, Large strongyles
S. equinus
Trichonema spp
Thodonomerus spp
Parascaris equorum
Oxyuris spp
Ascarids
 Small strongyles

DIRECTIONS FOR USE:
 Shake container well before using this product.

ADMINISTRATION:

WSD FENBENDAZOLE 100 can be administered orally through standard accurate drenching equipment. Clean equipment thoroughly before and after use.

DOSE RATES:
 Dose rates should be based on the heaviest animal by live weight in each group (bulls, cows, steers and calves), horses) and these should be weighed with scales or weighband. Do not underdose. A representative sample of animals should be weighed before treatment either with scales or weighband. Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Drenching into two or more bins may be appropriate, to avoid excessive overdosing.

CATTLE: 3.75ml/50kg body weight (7.5mg Fenbendazole per kg body weight)
HORSES: 1ml/10kg body weight (10mg Fenbendazole per kg body weight)
 For the control of *Strongyloides westeri* in foals: 5ml/10kg body weight (50mg Fenbendazole per kg body weight)

Bodyweight	Dose Rate	10L Treats	5L Treats	1L Treats
50kg	3.75ml	266	133	26
100kg	7.5ml	533	266	53
150kg	11.25ml	800	400	80
200kg	15.0ml	1066	533	107
250kg	18.75ml	1333	666	133
300kg	22.5ml	1600	800	160
350kg	26.25ml	1866	933	187
400kg	30.0ml	2133	1066	213
450kg	33.75ml	2400	1200	240
500kg	37.5ml	2666	1333	267
550kg	41.25ml	2933	1466	293
600kg	45.0ml	3200	1600	320
650kg	48.75ml	3466	1733	347
700kg	52.5ml	3733	1866	373
750kg	56.25ml	4000	2000	400
Over 750kg	3.75ml per 50kg	177	88	17

Bodyweight	Dose Rate	10L Treats	5L Treats	1L Treats
100kg	10ml	1000	500	100
200kg	20ml	2000	1000	200
250kg	25ml	2500	1250	250
300kg	30ml	3000	1500	300
350kg	35ml	3500	1750	350
400kg	40ml	4000	2000	400
450kg	45ml	4500	2250	450
500kg	50ml	5000	2500	500
550kg	55ml	5500	2750	550
600kg	60ml	6000	3000	600
650kg	65ml	6500	3250	650
700kg	70ml	7000	3500	700
750kg	75ml	7500	3750	750
Over 750kg	1ml per 10kg	133	67	13

DISPOSAL:

Do not dispose of undiluted chemicals on-site. Break, crush, puncture or bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roads. Empty containers and product should not be burnt.

STORAGE:

Store below 30°C (room temperature) in the closed, original container in a dry, well-ventilated area, and protect from light.

CONDITIONS OF SALE

All Western Stock Distributors products are of a high quality, being subjected to rigid quality control. The manufacturer, however, is unable to control the product's subsequent handling or use or the circumstances of each handling or use. This product is sold on the express condition that neither the manufacturer nor the distributor is under any liability for any loss or damage that may arise through such handling or use. If these conditions are unacceptable the container should be returned unopened for credit.

WESTERN STOCK DISTRIBUTORS

A.B.N. 58 009 372 525
 Koopjan Avenue, South Guildford WA 6055

NRA Approval No. 40427/1100

Batch No.:

Expiry: