



SPECIFICATION SHEET

WSD SPURT

Product Description

WSD SPURT is a treatment for body lice of sheep.

Type of Preparation

WSD SPURT is a pour-on formulation for topical use containing 25g/L **Cypermethrin**- a synthetic pyrethroid insecticide- and a scourable dye.

Indications

Sheep and lambs:

Control of biting lice (*Bovicola ovis*) in sheep off-shears

Milking sheep: Do NOT use in sheep producing milk for human consumption

Dosage

Always refer to the label. Applied at a dose rate of 1ml/5Kg body weight. The correct dose for the weight range of sheep should be applied as a single continuous strip 20-40cm in length down the middle of the back within 24 hours of shearing.

It is recommended that treated sheep be held together in the yards for 2 hours after treatments and not be let straight out of the race into the paddock.

Advice on use

- **WSD SPURT** is applied undiluted.
- It must not be stored in any container other than the original including overnight. Do not leave **WSD SPURT** containers uncovered or in direct sunlight.
- It is critical that sheep be cleanly shorn without residual clumps or tufts of wool and **WSD SPURT** must be applied less than 24 hours after shearing to ensure effectiveness.
- Correct application is critical and it should not be applied as a single spot or to sheep which are wet.
- Light to moderate rainfall soon after application should not reduce efficacy.
- A clean muster and single shearing with immediate correct treatment of every animal is ideal for achieving effective control of lice.
- Lambs at foot cannot be treated, and therefore it is recommended that ewes due to lamb within 6 weeks or ewes with lambs at foot should not be treated with **WSD SPURT**.
- If a split shearing is inevitable it is critical that treated and untreated sheep be kept separate.
- The repeated use of any synthetic pyrethroid is likely to result in the emergence of resistance in sheep lice. It is therefore recommended to regularly alternate treatment with a dip such as **WSD FLOCKMASTER MKII**.
- Resistance to synthetic pyrethroids is present in some populations of sheep lice, and treatment failures may be due to resistant lice being present.

Withholding periods

Meat withholding: 3 days

Export slaughter interval: 14 days

Milk withholding: Not for use in sheep producing milk for human consumption

Wool withholding: Because this is an off-shears treatment, no wool WHP is established.

Shelf Life

24 months from the date of manufacture

Sizes Available

5 litre Backpack, 20 litre Drum



WSD SPURT

GENERAL INSTRUCTIONS

Shake or stir contents before use.

APPLICATION

For maximum benefit and efficiency, pour-ons need to be applied correctly in accordance with their instructions.

- a) Prior to the start of shearing, check your SPURT applicator for correct operation and calibration.
- b) Apply SPURT to shorn sheep up to 24 hours of shears using the correct and full dose. Control is achieved in 5-6 weeks.
- c) Apply the SPURT dose as a single backline strip (40mm to 60mm wide) along the centre of back – start at the shoulder and finish at the rump (20cm to 40cm).
- d) Ensure all shorn, treated sheep go into a cleanly mustered paddock, and are not mixed with untreated sheep.

Applied correctly, SPURT puts a full strength dose of insecticide on every sheep and works well even under such adverse conditions as wet weather or wet sheep. A scourable blue marker dye is included in the product for specific identification of treated animals.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than three days before slaughter for human consumption.

MILK: DO NOT USE on female sheep which are producing or may in the future produce milk or milk products for human consumption.

SAFETY DIRECTIONS

Product is poisonous if swallowed. It will irritate the eyes and skin. Facial skin contact may cause temporary facial numbness. Sensitive workers should use protective clothing. Avoid contact with eyes and skin. Wear elbow length PVC gloves. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone 13 11 26).

MATERIAL SAFETY DATA SHEET (MSDS)

Additional information is listed in the Material Safety Data Sheet, which can be obtained from the supplier.

PROTECTION OF FISH, CRUSTACEANS AND THE ENVIRONMENT

Dangerous to fish and bees. Do not contaminate streams, rivers and waterways with the product or used containers.

DISPOSAL

Do not dispose of undiluted chemicals on-site. Break, crush, puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt.

STORAGE

Store below 30°C (room temperature) in the closed, original container in a dry, well-ventilated area, and protect from sunlight.

CONDITIONS OF SALE:

All Western Stock Distributors products are of a high quality, being subjected to rigid quality control. The manufacturer, however, is unable to control the product's subsequent handling or use or the circumstances of each handling or use. This product is sold on the express condition that neither the manufacturer nor the distributor is under any liability for any loss or damage that may arise through such handling or use. If these conditions are unacceptable the container should be returned unopened for credit.

WESTERN STOCK DISTRIBUTORS

A.B.N. 58 009 372 525
Kooljar Avenue, South Guildford WA 6055
Telephone: (08) 9277 5013

APVMA Approval No: 39065/1200

Batch No:

Expiry:

Sheep	Dose	1L Pack Treats	5L Pack Treats	20L Pack Treats
Under 10kg	2mL	500	2500	10000
11-20kg	4mL	250	1250	5000
21-30kg	6mL	166	833	3333
31-40kg	8mL	125	625	2500
41-50kg	10mL	100	500	2000
51-60kg	12mL	83	416	1666
61-70kg	14mL	71	357	1428
71-75kg	15mL	66	333	1333

Sheep in excess of 75kg dose at 1mL per 5kg

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.



DOSAGE
Dose the mob to the heaviest animal by liveweight in each group (ewes, wethers, rams and lambs).
A representative sample of animals should be weighed before treatment either with scales or weighband. Where there is a large variation in size within the group, dose rates should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate to avoid excessive overdosing.

RESTRAINTS
Must only be used within 24 hours of shearing.
Do not treat ewes less than 6 weeks before lambing commences, as live lice can persist and infect the lambs.
Do not treat lambing ewes or ewes with lambs at foot.
Do not mix treated sheep with untreated sheep until 6 weeks after treatment.

Do not use on sheep which have not been cleanly shorn.
DO NOT USE on female sheep which are producing or may in the future produce milk or milk products for human consumption.

OFF-SHEARS, POUR-ON SHEEP LICE CONTROL
Contains Cypermethrin and is effective against pyrethroid susceptible strains of *Gamulha ovis*.

DIRECTIONS FOR USE

RESTRAINTS

Must only be used within 24 hours of shearing.
Do not treat ewes less than 6 weeks before lambing commences, as live lice can persist and infect the lambs.
Do not treat lambing ewes or ewes with lambs at foot.
Do not mix treated sheep with untreated sheep until 6 weeks after treatment.

Do not use on sheep which have not been cleanly shorn.
DO NOT USE on female sheep which are producing or may in the future produce milk or milk products for human consumption.

DOSAGE

Dose the mob to the heaviest animal by liveweight in each group (ewes, wethers, rams and lambs).
A representative sample of animals should be weighed before treatment either with scales or weighband. Where there is a large variation in size within the group, dose rates should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate to avoid excessive overdosing.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than three days before slaughter for human consumption.

MILK: DO NOT USE on female sheep which are producing or may in the future produce milk or milk products for human consumption.

SAFETY DIRECTIONS

Product is poisonous if swallowed. It will irritate the eyes and skin. Facial skin contact may cause temporary facial numbness. Sensitive workers should use protective clothing. Avoid contact with eyes and skin. Wear elbow length PVC gloves. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone 13 11 26).

MATERIAL SAFETY DATA SHEET (MSDS)

Additional information is listed in the Material Safety Data Sheet, which can be obtained from the supplier.